

The shell is conical and can measure up to 30 cm (12") but usually ranges from 5-13 cm (2-5"). It usually has more than seven whorls (Fig. 2).



Image: Florida Dept. of Agriculture and Consumer Services.

Fig.2. Shell of *A. fulica*.

The colour of the shell varies with diet and the environment. The shell is usually brown with reddish-brown and light yellow or cream streaks (Fig. 3).



Image: Roberta Zimmerman, USDA APHIS

Fig. 3. *A. fulica* adults.

Where do I look?

The giant African snail is nocturnal, but may become active at dawn and dusk if it is very wet. They prefer damp, shady places and avoid direct sunlight. In the day they are commonly found:

- under ledges, bricks, rocks, logs, plant mats, decaying leaves, or discarded containers; and
- in or on plants, trees, heavy vegetation, brick holes, crevices.

What can I do?

1. Report sightings
 - call the nearest regional agricultural office.
 - email pictures to agriquarantine@gmail.com
2. Do not put live snail in the garbage.
3. Do not move soils/debris from infested areas.
4. Check your vehicles and machinery for snails.
5. Use gloves when handling snails.
6. Crush smaller snails.
7. Use slug snail bait as prescribed.
8. **Help stop the spread of the giant African Snail**



Giant African Snail

(*Achatina Fulica*)

**A notifiable pest in the
Commonwealth of Dominica**
(Act 10 of 1986)

Acknowledgement:
Plant & Protection Quarantine Services
Botanical Gardens
Roseau
Tel: 266-3820

Ministry of Agriculture & Forestry
Plant Protection & Quarantine Services



images: Chandrabhan Shrivast, Research Division, Ministry of Food, Production and



Marine Affairs, Trinidad and Tobago



image: Giant African land snail © Josef Dvořák
<http://ocassweb.guyana.gov.gy/photo/AF/Min/3/Anw/Ar/M/1/10/10>

What is it?

The giant African snail is one of the world's most destructive land snails. It is an alien invasive species of quarantine importance.

Where is it present?

The giant African snail is native to East Africa. It was introduced into Asia, the Pacific Region and Brazil. It was found numerous times in the USA and was eradicated. In the 1980's it was introduced into the Caribbean, and today it is present in Guadeloupe, Martinique, St Lucia, Barbados, Antigua and Dominica.

How is it spread?

Live giant African snails spread across the world through deliberate introduction, illegal trade and contaminated material. Also, dormant snails may be mistaken as empty shells and moved by collectors. Within a country, the snail spreads by hitching a ride on vehicles, shipping containers, machinery, crates, and plant material. Snail eggs can be spread by moving infested soil, garbage and plant debris.

What does it eat?

In its native range, the giant African snail is a scavenger found on decayed vegetation. As an invasive species it has been reported on over **500 plant types** including:

- vegetable crops (cabbage, ochro, cucumbers, pumpkin, melons, spinach, lettuce, tomato, pepper, broccoli, cauliflower, peas, beans, breadfruit);
- fruit trees (citrus, bananas, papaya, soursop);
- field crops (dasheen, cassava, yam, sweet potato, coffee, cocoa, sugarcane);
- forest trees (teak, mahogany); and
- ornamentals (bougainvillea, poinsettia, orchid, rose).

Why should I care?

The giant African snail is a serious pest of agriculture and tourism (damaging landscapes). It can cause significant economic losses. In Florida it was estimated that an annual loss of \$US11 million (in 1969) would have been incurred if its population had not been controlled.

The giant African Snail is a vector of the rat lungworm (*Angiostrongylus cantonensis*) that causes eosinophilic meningitis in humans. This parasite can be contracted by eating improperly cooked snail meat or by handling live snails and transferring snail mucus to our eyes, nose, and mouth.

The giant African Snail can live up to 10 years and has the ability to become dormant in unfavourable

conditions. Each snail carries both male and female sex organs. After mating both snails produce eggs. A snail can produce up to 500 eggs in one laying, and 1200 individuals in a year, thereby establishing a very large population in a relatively short time. **Once this snail is established, eradication is difficult and expensive.**

What do I look for?

The giant African snail is large, weighing as much as 1 kg. It lays eggs in batches of 100-500. Eggs are round to oval, about 5 mm (1/4") in length and cream to yellow (Fig. 1).



Image: Yuri Yashin, *achatina.ru*, Bugwood.org

Fig.1. A clutch of *A. fulica* eggs.